

Integrable string models with constant torsion in terms of chiral invariants of $SU(n)$, $SO(n)$, $SP(n)$ groups

V.D Gershun¹

¹*ITP, NSC Kharkov Institute of Physics and Technology, Kharkov, Ukraine*

We used the invariant local chiral currents of principal chiral models for $SU(n)$, $SO(n)$, $SP(n)$ groups to construct new integrable string equations of hydrodynamic type on the Riemann space of the chiral primitive invariant currents and on the chiral non-primitive Casimir operators as Hamiltonians.

PACS numbers: 02.20.Sv, 02.40.Vv, 11.17.-w, 11.30.Rd, 21.60.Fw

1. INTRODUCTION

For quantum description of string model we must to have classical solutions of the string in the background fields. String theory in suitable space-time backgrounds can be considered as principal chiral model. The integrability of the classical principal chiral model is manifested through an infinite set of conserved charges, which can form non-abelian algebra. Any charge from the commuting subset of charges and any Casimir operators of charge algebra can be considered as Hamiltonian. The polynomials of local chiral currents was considered by Goldshmidt and Witten [1] (see also [2]). The local conserved chiral charges in principal chiral models was considered by Evans, Hassan, MacKay, Mountain [5]. The integrable string models of hydrodynamic type was considered by author [3], [4]. In section 2 the author obtained the basis local invariant chiral currents for $SU(n)$ which they form closed algebra under Poisson brackets of hydrodynamic type. The author constructed new invariant symmetric tensors and obtained relations between new invariant tensors and basis tensors. In section 3 the author obtained new integrable string equations for $SU(3)$ group for chiral currents of $SU(3)$ group as local fields of Riemann space with non-primitive chiral Casimirs as Hamiltonians. The non-primitive invariant chiral charges for $SU(n)$ $n \geq 4$ are not Casimirs and they can not used as Hamiltonians to construct new integrable equations for this groups. The author show that new integrable string equations do not arise for $SO(2l+1)$, $SO(2l)$, $SP(2l)$ groups for $l \geq 2$.

A string model is described by the Lagrangian

$$L = \frac{1}{2} \int_0^{2\pi} \eta^{\alpha\beta} g_{ab}(\phi(t, x)) \frac{\partial \phi^a(t, x)}{\partial x^\alpha} \frac{\partial \phi^b(t, x)}{\partial x^\beta} dx$$

and by two first kind constraints:

$$g_{ab}(\phi(x)) \left[\frac{\partial \phi^a(x)}{\partial t} \frac{\partial \phi^b(x)}{\partial t} + \frac{\partial \phi^a(x)}{\partial x} \frac{\partial \phi^b(x)}{\partial x} \right] \approx 0,$$

$$g_{ab}(\phi(x)) \frac{\partial \phi^a(x)}{\partial t} \frac{\partial \phi^b(x)}{\partial x} \approx 0.$$

The target space local coordinates $\phi^a(x)$, $a = 1, \dots, n$ belong to certain given smooth n -dimensional manifold M^n with nondegenerate metric tensor

$$g_{ab}(\phi(x)) = \eta_{\mu\nu} e_a^\mu(\phi(x)) e_b^\nu(\phi(x)),$$

where $\mu, \nu = 1, \dots, n$ are indexes of tangent space to manifold M^n on some point $\phi^a(x)$. The veilbein $e_a^\mu(\phi)$ and its inverse $e_\mu^a(\phi)$ satisfy to the conditions:

$$e_a^\mu e_\mu^b = \delta_a^b, \quad e_a^\mu e^\nu = \eta^{\mu\nu}.$$

The coordinates x^α , $x^0 = t$, $x^1 = x$ belong to world sheet with metric tensor $g_{\alpha\beta}$ in conformal gauge. The string equations of motion have the form:

$$\eta^{\alpha\beta} [\partial_{\alpha\beta} \phi^a + \Gamma_{bc}^a(\phi) \partial_\alpha \phi^b \partial_\beta \phi^c] = 0, \quad \partial_\alpha = \frac{\partial}{\partial x^\alpha}, \quad \alpha = 0, 1,$$

where

$$\Gamma_{bc}^a(\phi) = \frac{1}{2} e_\mu^a \left[\frac{\partial e_b^\mu}{\partial \phi^c} + \frac{\partial e_c^\mu}{\partial \phi^b} \right]$$

is connection.

The Hamiltonian has form:

$$H = \frac{1}{2} \int_0^{2\pi} [\eta^{\mu\nu} J_{0\mu} J_{0\nu} + \eta_{\mu\nu} J_1^\mu J_1^\nu] dx,$$

where $J_{0\mu}(\phi) = e_\mu^a(\phi) p_a$, $J_1^\mu(\phi) = e_a^\mu \frac{\partial}{\partial x} \phi^a$ and $p_a(t, x) = \eta_{\mu\nu} e_a^\mu e_b^\nu \frac{\partial}{\partial t} \phi^b$ is canonical momentum.

Let us introduce chiral currents:

$$U^\mu = \eta^{\mu\nu} J_{0\nu} + J_1^\mu, \quad V^\mu = \eta^{\mu\nu} J_{0\nu} - J_1^\mu$$

The commutation relations of chiral currents U^μ are following:

$$\{U^\mu(x), U^\nu(y)\} = C_\lambda^{\mu\nu} [\frac{3}{2}U^\lambda(x) - \frac{1}{2}V^\lambda(x)]\delta(x-y) - \eta^{\mu\nu} \frac{\partial}{\partial x} \delta(x-y), \quad (1)$$

where

$$C_{\nu\lambda}^\mu(\phi) = e_\nu^a e_\lambda^b [\frac{\partial e_a^\mu}{\partial \phi^b} - \frac{\partial e_b^\mu}{\partial \phi^a}]$$

is torsion. Equations of motion in light-cone coordinates

$$x^\pm = \frac{1}{2}(t \pm x), \quad \frac{\partial}{\partial x^\pm} = \frac{\partial}{\partial t} \pm \frac{\partial}{\partial x}$$

have form:

$$\partial_- U^\mu = C_{\nu\lambda}^\mu(\phi(x)) U^\nu V^\lambda, \quad \partial_- V^\mu = C_{\nu\lambda}^\mu(\phi(x)) V^\nu U^\lambda.$$

In the case of null torsion:

$$C_{\nu\lambda}^\mu = 0, \quad e_a^\mu(\phi) = \frac{\partial e^\mu}{\partial \phi^a}, \quad \Gamma_{bc}^a(\phi) = e_\mu^a \frac{\partial^2 e^\mu}{\partial \phi^b \partial \phi^c}, \quad R_{\nu\lambda\rho}^\mu(\phi) = 0$$

string model is integrable one. The Hamiltonian equations of motion under Hamiltonian H are described two independent left and right movers: $U^\mu(t+x)$ and $V^\mu(t-x)$.

2. INTEGRABLE STRING MODELS WITH CONSTANT TORSION

Let us come back to commutations relations of chiral currents. Let torsion $C_{\nu\lambda}^\mu(\phi(x)) \neq 0$ and $C_{\mu\nu\lambda} = f_{\mu\nu\lambda}$ are structure constant of simple Lie algebra. We will consider string model with constant torsion in light-cone gauge in target space. This model coincides to principal chiral model on compact simple Lie group. We can not divide motion on right and left mover because of chiral currents $\partial_- U^\mu = f_{\nu\lambda}^\mu U^\nu V^\lambda$, $\partial_- V^\mu = f_{\nu\lambda}^\mu V^\nu U^\lambda$ are not conserve. The correspondent charges are not Casimirs. Evans, Hassan, MacKay, Mountain constructed local invariant chiral currents as polynomials of initial chiral currents of $SU(n)$, $SO(n)$, $SP(n)$ principal chiral models and they found such combination of them, that corresponding charges are Casimir operators of this dynamical systems. Their paper was based on the paper of de Azcarraga, Macfarlane, MacKay, Perez Bueno [6] about invariant tensors for simple Lie algebras. Let t_μ are $n \otimes n$ traceless hermitian matrix representations of generators Lie algebra:

$$[t_\mu, t_\nu] = 2if_{\mu\nu\lambda} t_\lambda, \quad Tr(t_\mu t_\nu) = 2\delta_{\mu\nu}.$$

Here is additional relation for $SU(n)$ algebra:

$$\{t_\mu, t_\nu\} = \frac{4}{n}\delta_{\mu\nu} + 2d_{\mu\nu\lambda}t_\lambda, \quad \mu = 1, \dots, n^2 - 1.$$

De Azcarraga et. al. gave some examples of invariant tensors of simple Lie algebras and they gave general method to calculate them. Invariant tensors may to construct as invariant symmetric polynomials on $SU(n)$:

$$d_M = d_{(\mu_1 \dots \mu_M)} = \frac{1}{M!} STr(t_{\mu_1} \dots t_{\mu_M}),$$

where STr means of completely symmetrized product of matrices and $d_{(\mu_1 \dots \mu_M)}$ is totally symmetric tensor and $M = 2, 3, \dots, \infty$. Another family of invariant symmetric tensors, such named d -family, based on the product of the symmetric structure constant $d_{\mu\nu\lambda}$ of $SU(n)$ algebra:

$$C_M = d_{(\mu_1 \dots \mu_M)} = d_{(\mu_1 \mu_2}^{k_1} d_{\mu_3}^{k_1 k_2} \dots d_{\mu_{M-2}}^{k_{M-2} k_{M-3}} d_{\mu_{M-1} \mu_M}^{k_{M-3}},$$

where $C_3 = d_{\mu\nu\lambda}$ and $M = 4, 5, \dots, \infty$.

Here are $n - 1$ primitive invariant tensors on $SU(n)$. The invariant tensors for $M \geq n$ are functions of primitive tensors. Evans et.al. introduced local chiral currents based on the invariant symmetric polynomials on simple Lie groups:

$$J_M(U) = d_{\mu_1 \dots \mu_M} U^{\mu_1} \dots U^{\mu_M},$$

where $U = t_\mu U^\mu$ and $\mu = 1, \dots, n^2 - 1$. It is possible to compose the invariant symmetric polynomials $J_M(U)$ to basis invariant tensors $C_M(U)$:

$$C_2(U) = \eta_{\mu\nu} U^\mu U^\nu, \quad C_3(U) = d_{\mu\nu\lambda} U^\mu U^\nu U^\lambda, \quad C_M(U) = d_{\mu_1 \mu_2}^{k_1} d_{\mu_3}^{k_1 k_2} \dots d_{\mu_{M-1} \mu_M}^{k_{M-3}} U_{\mu_1} U_{\mu_2} \dots U_{\mu_M}, \quad (2)$$

where $M = 4, 5, \dots, \infty$. The author obtained following expression for local invariant chiral currents $J_M(U)$:

$$\begin{aligned} J_2 &= 2C_2, \quad J_3 = 2C_3, \quad J_4 = 2C_4 + \frac{4}{n}C_2^2, \quad J_5 = 2C_5 + \frac{8}{n}C_2C_3, \\ J_6 &= 2C_6 + \frac{4}{n}C_3^2 + \frac{8}{n}C_2C_4 + \frac{8}{n^2}C_2^3, \\ J_7 &= 2C_7 + \frac{8}{n}C_3C_4 + \frac{8}{n}C_2C_5 + \frac{24}{n^2}C_2^2C_3, \\ J_8 &= 2C_8 + \frac{4}{n}C_4^2 + \frac{8}{n}C_3C_5 + \frac{8}{n}C_2C_6 + \frac{24}{n^2}C_2C_3^2 + \frac{24}{n^2}C_2^2C_4 + \frac{16}{n^3}C_2^4, \end{aligned} \quad (3)$$

$$J_9 = 2C_9 + \frac{8}{n}C_4C_5 + \frac{8}{n}C_3C_6 + \frac{8}{n}C_2C_7 + \frac{8}{n^2}C_3^3 + \frac{48}{n^2}C_3C_4 + \frac{24}{n^2}C_2^2C_5 + \frac{64}{n^3}C_2^3C_3.$$

The commutation relations of invariant chiral currents $J_M(U(x))$ show, that these currents are not densities of dynamical Casimir operators. We considered the basis family of invariant chiral currents $C_M(U)$ and we proved that invariant chiral currents $C_M(U)$ forms closed algebra under canonical PB and corresponding charges are dynamical Casimir operators. The commutation relations of invariant chiral currents $C_M(U(x))$ and $C_N(U(y))$ for $M, N = 2, 3, 4$ and for $M = 2, N = 2, 3, \dots, \infty$ are the following:

$$\{C_M(x), C_N(y)\} = -MNC_{M+N-2}(x) \frac{\partial}{\partial x} \delta(x-y) - \frac{MN(N-1)}{M+N-2} \frac{\partial C_{M+N-2}(x)}{\partial x} \delta(x-y).$$

The commutations relations for $M \geq 5, N \geq 3$ are following (we show one formula for C_8 on the right side as example only):


$$\begin{aligned} \{C_5(x), C_3(y)\} &= -[12C_6(x) + 3C_{6,1}(x)] \frac{\partial}{\partial x} \delta(x-y) - \frac{1}{3} \frac{\partial}{\partial x} [12C_6(x) + 3C_{6,1}(x)] \delta(x-y), \\ \{C_5(x), C_4(y)\} &= -[16C_7(x) + 4C_{7,1}(x)] \frac{\partial}{\partial x} \delta(x-y) - \frac{3}{7} \frac{\partial}{\partial x} [16C_7(x) + 4C_{7,1}(x)] \delta(x-y), \\ \{C_6(x), C_3(y)\} &= -[12C_7(x) + 6C_{7,1}(x)] \frac{\partial}{\partial x} \delta(x-y) - \frac{2}{7} \frac{\partial}{\partial x} [12C_7(x) + 6C_{7,1}(x)] \delta(x-y), \\ \{C_5(x), C_5(y)\} &= -[16C_8(x) + 8C_{8,1}(x) + C_{8,2}(x)] \frac{\partial}{\partial x} \delta(x-y) - \\ &\quad \frac{1}{2} \frac{\partial}{\partial x} [16C_8(x) + 8C_{8,1}(x) + C_{8,2}(x)] \delta(x-y), \\ \{C_6(x), C_4(y)\} &= -[16C_8(x) + 8C_{8,3}(x)] \frac{\partial}{\partial x} \delta(x-y) - \\ &\quad \frac{3}{8} \frac{\partial}{\partial x} [16C_8(x) + 8C_{8,3}(x)] \delta(x-y), \\ \{C_7(x), C_3(y)\} &= -[12C_8(x) + 6C_{8,1}(x) + 3C_{8,3}(x)] \frac{\partial}{\partial x} \delta(x-y) - \\ &\quad \frac{1}{4} \frac{\partial}{\partial x} [12C_8(x) + 6C_{8,1}(x) + 3C_{8,3}(x)] \delta(x-y), \\ \{C_8(x), C_3(y)\} &= -[12C_9(x) + 6C_{9,1}(x) + 6C_{9,2}(x)] \frac{\partial}{\partial x} \delta(x-y) - \\ &\quad \frac{2}{9} \frac{\partial}{\partial x} [12C_9(x) + 6C_{9,1}(x) + 6C_{9,2}(x)] \delta(x-y), \\ \{C_7(x), C_4(y)\} &= -[16C_9(x) + 8C_{9,2}(x) + 4C_{9,3}(x)] \frac{\partial}{\partial x} \delta(x-y) - \\ &\quad \frac{1}{3} \frac{\partial}{\partial x} [16C_9(x) + 8C_{9,2}(x) + 4C_{9,3}(x)] \delta(x-y), \end{aligned} \tag{4}$$

$$\begin{aligned}
\{C_6(x), C_5(y)\} &= -[16C_9(x) + 4C_{9,1}(x) + 8C_{9,2}(x) + 2C_{9,4}(x)]\frac{\partial}{\partial x}\delta(x-y) - \\
&\quad \frac{4}{9}\frac{\partial}{\partial x}[16C_9(x) + 4C_{9,1}(x) + 8C_{9,2}(x) + 2C_{9,4}(x)]\delta(x-y), \\
\{C_9(x), C_3(y)\} &= -[12C_{10}(x) + 6C_{10,1}(x) + 6C_{10,2}(x) + 3C_{10,3}(x)]\frac{\partial}{\partial x}\delta(x-y) - \\
&\quad \frac{1}{5}\frac{\partial}{\partial x}[12C_{10}(x) + 6C_{10,1}(x) + 6C_{10,2}(x) + 3C_{10,3}(x)]\delta(x-y), \\
\{C_8(x), C_4(y)\} &= -[16C_{10}(x) + 8C_{10,2}(x) + 8C_{10,4}(x)]\frac{\partial}{\partial x}\delta(x-y) - \\
&\quad \frac{3}{10}\frac{\partial}{\partial x}[16C_{10}(x) + 8C_{10,2}(x) + 8C_{10,4}(x)]\delta(x-y), \\
\{C_7(x), C_5(y)\} &= -[16C_{10}(x) + 8C_{10,3}(x) + 4C_{10,1}(x) + 4C_{10,4}(x) + 2C_{10,5}(x) + C_{10,6}(x)]\frac{\partial}{\partial x}\delta(x-y) - \\
&\quad \frac{2}{5}\frac{\partial}{\partial x}[16C_{10}(x) + 8C_{10,3}(x) + 4C_{10,1}(x) + 4C_{10,4}(x) + 2C_{10,5}(x) + C_{10,6}(x)]\delta(x-y), \\
\{C_6(x), C_6(y)\} &= -[16C_{10}(x) + 16C_{10,2}(x) + 4C_{10,7}(x)]\frac{\partial}{\partial x}\delta(x-y) - \\
&\quad \frac{1}{2}\frac{\partial}{\partial x}[16C_{10}(x) + 16C_{10,2}(x) + 4C_{10,7}(x)]\delta(x-y).
\end{aligned}$$

The new dependent invariant chiral currents $C_{6,1}$, $C_{7,1}$, $C_{8,1} - C_{8,3}$, $C_{9,1} - C_{9,4}$, $C_{10,1} - C_{10,7}$ have the similar form (we show formulas for C_8 on the right side as example only):

$$\begin{aligned}
C_{8,1} &= [d_{\mu\nu}^k d_{\lambda}^{kl} d_{\rho}^{ln}][d_{\sigma\varphi}^m][d_{\tau\theta}^p]d^{nmp}(U^8)^{\mu\nu\lambda\rho\sigma\varphi\tau\theta}, \\
C_{8,2} &= [d_{\mu\nu}^k][d_{\lambda\rho}^l][d_{\sigma\varphi}^n][d_{\tau\theta}^m]d^{klp}d^{nmp}(U^8)^{\mu\nu\lambda\rho\sigma\varphi\tau\theta}. \\
C_{8,3} &= [d_{\mu\nu}^k d_{\lambda}^{kl}][d_{\rho\sigma}^m d_{\varphi}^{nm}][d_{\tau\theta}^p]d^{lmp}(U^8)^{\mu\nu\lambda\rho\sigma\varphi\tau\theta},
\end{aligned}$$

The following picture show the graphic image of invariant chiral currents $C_M(U)$ and the difference between C_8 and $C_{8,1} - C_{8,3}$.



Let us note that these PB's (4) are PB's of hydrodynamic type. The ultra local term with antisymmetric structure constant $f_{\mu\nu\lambda}$ in commutation relation of chiral currents U^μ

(1) does not contribution to commutation relations of invariant chiral currents because of totally symmetric invariant tensors $d_{(\mu_1 \dots \mu_M)}$. Therefore chiral currents $C_M(U(x))$ form closed algebra under canonical PB.

The new dependent invariant chiral currents and the new dependent totally symmetric invariant tensors for $SU(N)$ group can be obtained under different order of calculation of trace of the product of the generators of $SU(n)$ algebra. Let us mark the matrix product of two generators t_μ, t_ν in round brackets:

$$(t_\mu t_\nu) = \frac{2}{n} \delta_{\mu\nu} + (d_{\mu\nu}^k + i f_{\mu\nu}^k) t_k.$$

The expression of invariant chiral currents $J_M(U)$ depends of the order of the matrix product of two generators in general list of generators. For example:

$$\begin{aligned} J_8 &= Tr[t(\underline{tt})t(\underline{tt})t] = 2C_8 + \frac{4}{n}C_4^2 + \frac{8}{n}C_3C_5 + \frac{8}{n}C_2C_6 + \frac{24}{n^2}C_2C_3^2 + \frac{24}{n^2}C_2^2C_4 + \frac{16}{n^3}C_2^4, \\ J_8 &= Tr[(\underline{tt})(\underline{tt})t(\underline{tt})t] = 2C_{8,1} + \frac{4}{n}C_4^2 + \frac{4}{n}C_3C_5 + \frac{24}{n^2}C_2C_3^2 + \frac{12}{n}C_2C_6 + \frac{24}{n^2}C_2^2C_4 + \frac{16}{n^3}C_2^4, \\ J_8 &= Tr[(\underline{tt})(\underline{tt})(\underline{tt})(\underline{tt})] = 2C_{8,2} + \frac{4}{n}C_4^2 + \frac{16}{n}C_2C_{6,1} + \frac{32}{n^2}C_2^2C_4 + \frac{16}{n^3}C_2^4, \\ J_8 &= Tr[t(\underline{tt})(\underline{tt})(\underline{tt})t] = 2C_{8,3} + \frac{12}{n}C_2C_6 + \frac{8}{n}C_3C_5 + \frac{24}{n^2}C_2^2C_4 + \frac{24}{n^2}C_2C_3^2 + \frac{16}{n}C_2^4, \\ J_9 &= Tr[t(\underline{tt})t(\underline{tt})t(\underline{tt})t] = 2C_9 + \frac{8}{n}C_4C_5 + \frac{8}{n}C_3C_6 + \frac{8}{n}C_2C_7 + \frac{8}{n^2}C_3^3 + \frac{48}{n^2}C_2C_3C_4 + \frac{24}{n^2}C_2^2C_5 + \frac{64}{n^3}C_2^3C_3, \\ J_9 &= Tr[t(\underline{tt})t(\underline{tt})(\underline{tt})] = \\ &\left\{ \begin{aligned} &2C_{9,1} + \frac{4}{n}C_4C_5 + \frac{4}{n}C_2C_7 + \frac{4}{n}C_2C_{7,1} + \frac{8}{n}C_3C_{6,1} + \frac{32}{n^2}C_2C_3C_4 + \frac{32}{n^2}C_2^2C_5 + \frac{64}{n^3}C_2^3C_3 \\ &2C_{9,4} + \frac{4}{n}C_2C_7 + \frac{4}{n}C_2C_{7,1} + \frac{12}{n}C_3C_{6,1} + \frac{32}{n^2}C_2C_3C_4 + \frac{32}{n^2}C_2^2C_5 + \frac{64}{n^3}C_2^3C_3, \end{aligned} \right. \\ J_9 &= Tr[t(\underline{tt})t(\underline{tt})t(\underline{tt})] = \\ &\left\{ \begin{aligned} &2C_{9,2} + \frac{4}{n}C_4C_5 + \frac{8}{n}C_3C_6 + \frac{8}{n}C_2C_7 + \frac{4}{n}C_2C_{7,1} + \frac{8}{n^2}C_3^3 + \frac{40}{n^2}C_2C_3C_4 + \frac{32}{n^2}C_2^2C_5 + \frac{64}{n^3}C_2^3C_3, \\ &2C_{9,3} + \frac{8}{n}C_2C_7 + \frac{4}{n}C_2C_{7,1} + \frac{12}{n}C_3C_6 + \frac{8}{n}C_3^3 + \frac{40}{n^2}C_2C_3C_4 + \frac{32}{n^2}C_2^2C_5 + \frac{64}{n^3}C_2^3C_3, \end{aligned} \right. \end{aligned}$$

where $t = t_\mu U^\mu$ and two variants of two last expressions for $J_9(U)$ obtained from two variants of expression for $J_6(U)$ during calculation $J_9(U)$. Because result of calculation does not depend of order of calculation, we can obtain relations between new invariant chiral currents and basis invariant currents $C_M(U)$:

$$C_{6,1} = C_6 + \frac{2}{n}C_3^2 - \frac{2}{n}C_2C_4,$$

$$\begin{aligned}
C_{7,1} &= C_7 + \frac{4}{n}C_3C_4 - \frac{4}{n}C_2C_5, \\
C_{8,1} &= C_8 + \frac{2}{n}C_3C_5 - \frac{2}{n}C_2C_6, \\
C_{8,2} &= C_8 + \frac{4}{n}C_3C_5 - \frac{4}{n}C_2C_6 - \frac{4}{n^2}C_2C_3^2 + \frac{4}{n^2}C_2^2C_4, \\
C_{8,3} &= C_8 + \frac{2}{n}C_4^2 - \frac{2}{n}C_2C_6, \\
C_{9,1} &= C_9 + \frac{2}{n}C_4C_5 - \frac{4}{n^2}C_3^3 + \frac{8}{n^2}C_2C_3C_4 + \frac{4}{n^2}C_2^2C_5, \\
C_{9,2} &= C_9 + \frac{2}{n}C_4C_5 - \frac{2}{n}C_2C_7 - \frac{4}{n^2}C_2C_3C_4 + \frac{4}{n^2}C_2^2C_5, \\
C_{9,3} &= C_9 + \frac{4}{n}C_4C_5 - \frac{2}{n}C_2C_7 - \frac{2}{n}C_3C_6 - \frac{4}{n^2}C_2C_3C_4 + \frac{4}{n^2}C_2^2C_5, \\
C_{9,4} &= C_9 + \frac{4}{n}C_4C_5 - \frac{2}{n}C_3C_6 - \frac{8}{n^2}C_3^3 + \frac{12}{n^2}C_2C_3C_4 + \frac{4}{n^2}C_2^2C_5.
\end{aligned} \tag{5}$$

Hence we can obtain the new relations for symmetric tensors:

$$\begin{aligned}
d_{(\mu\nu}^k d_{\lambda\rho}^l d_{\sigma\varphi}^n d_{\tau\theta}^{nm} d_{\theta}^{mp} d^{klp} &= d_{(\mu\nu}^k d_{\lambda}^{kl} d_{\rho}^{ln} d_{\sigma}^{nm} d_{\varphi}^{mp} d_{\tau\theta}^p) + \frac{4}{n} d_{(\mu\nu\lambda} d_{\rho\sigma}^k d_{\varphi}^{kl} d_{\tau\theta}^l) - \frac{2}{n} \delta_{(\mu\nu} d_{\lambda\rho}^k d_{\sigma}^{kl} d_{\varphi}^{ln} d_{\tau\theta}^n), \\
d_{(\mu\nu}^k d_{\lambda\rho}^l d_{\sigma\varphi}^n d_{\tau\theta}^{nm} d^{klp} d^{nmp} &= d_{(\mu\nu}^k d_{\lambda}^{kl} d_{\rho}^{ln} d_{\sigma}^{nm} d_{\varphi}^{mp} d_{\tau\theta}^p) + \frac{4}{n} d_{(\mu\nu\lambda} d_{\rho\sigma}^k d_{\varphi}^{kl} d_{\tau\theta}^l) - \\
&\quad - \frac{4}{n} \delta_{(\mu\nu} d_{\lambda\rho}^k d_{\sigma}^{kl} d_{\varphi}^{ln} d_{\tau\theta}^n) - \frac{4}{n^2} \delta_{(\mu\nu} d_{\lambda\rho\sigma} d_{\varphi\tau\theta}) + \frac{4}{n^2} \delta_{(\mu\nu} \delta_{\lambda\rho} d_{\sigma\varphi}^k d_{\tau\theta}^k).
\end{aligned}$$

The family of invariant chiral currents $C_M(U(x))$ satisfy to conservation equations $\partial_- C_M(U(x)) = 0$.

3. NEW INTEGRABLE STRING EQUATIONS

Let us apply hydrodynamic approach to integrable string models with constant torsion. In this case we must to consider the conserved primitive chiral currents currents $C_M(U(x))$, ($M = 2, 3, \dots, n-1$) as local fields of the Riemann manifold. The non-primitive local charges of invariant chiral currents with $M \geq n$ form the hierarchy of new Hamiltonians in bi-Hamiltonian approach to integrable systems. The commutation relations of invariant chiral currents are local PBs of hydrodynamic type.

The invariant chiral currents C_M with $M \geq 3$ for the $SU(3)$ group can be obtained from following relation:

$$d_{kln} d_{kmp} + d_{klm} d_{knp} + d_{klp} d_{knm} = \frac{1}{3} (\delta_{ln} \delta_{mp} + \delta_{lm} \delta_{np} + \delta_{lp} \delta_{nm}).$$

The corresponding invariant chiral currents for $SU(3)$ group have form:

$$C_{2N} = \frac{1}{3^{N-1}}(\eta_{\mu\nu}U^\mu U^\nu)^N = \frac{1}{3^{N-1}}(C_2)^N,$$

$$C_{2N+1} = \frac{1}{3^{N-1}}(\eta_{\mu\nu}U^\mu U^\nu)^{N-1}d_{klm}U^k U^l U^m = \frac{1}{3^{N-1}}(C_2)^{N-1}C_3.$$

The invariant chiral currents C_2, C_3 are local coordinates of the Riemann manifold M^2 . The local charges C_{2N} , $N \geq 2$ form hierarchy of Hamiltonians. The new nonlinear equations of motion for chiral currents are the following:

$$\frac{\partial f(x, t_N)}{\partial t_N} = -f^{N-1}(x)\frac{\partial f(x)}{\partial x}, \quad (6)$$

$$\frac{\partial g(x, t_N)}{\partial t_N} = -\frac{3}{2N-1}g(x)f^{N-2}(x)\frac{\partial f(x)}{\partial x} - \frac{1}{2N-1}f^{N-1}(x)\frac{\partial g(x)}{\partial x}.$$

Here $f(x) = C_2(U(x))$, $g(x) = C_3(U(x))$ and Hamiltonian has following form:

$$H_N(t_N) = \frac{1}{2N(2N-1)} \int_0^{2\pi} f^N(x, t_N) dx, \quad N = 2, \dots, \infty.$$

First equation for function $f(x, t_n)$ is generalized inviscid Burgers' equation. It has following solution:

$$f(x, t_N) = h[x - t_N f^{N-1}(x, t_N)] \quad (7)$$

and $h(x)$ is arbitrary function of x .

The construction of integrable equations with $SU(n)$ symmetries for $n \geq 4$ has difficulties of reduction non-primitive invariant currents to primitive currents. Following Balog, Feher, O'Raiheartaigh, Forgacs, Wipf we considered the generating function

$$A(x, \lambda) = \det(1 - \lambda U(x)) = \exp \text{Tr} \ln(1 - \lambda U^\mu(x) t_\mu).$$

We obtained following expressions for non-primitive invariant currents C_N :

$$SU(4) : C_5 \rightarrow \frac{2}{3}C_2C_3, \quad C_6 \rightarrow \frac{1}{6}C_3^2 + \frac{1}{2}C_2C_4, \quad C_7 \rightarrow \frac{1}{3}C_2^2C_3 + \frac{1}{6}C_3C_4,$$

$$C_8 \rightarrow \frac{7}{36}C_2C_3^2 + \frac{1}{4}C_2^2C_4, \quad C_9 \rightarrow \frac{1}{6}C_2^3C_3 + \frac{1}{36}C_3^3 + \frac{1}{6}C_2C_3C_4.$$

$$SU(5) : C_6 \rightarrow -\frac{3}{50}C_2^3 + \frac{4}{15}C_3^2 + \frac{7}{10}C_2C_4, \quad C_7 \rightarrow -\frac{3}{50}C_2^2C_3 + \frac{11}{30}C_3C_4 + \frac{3}{5}C_2C_5,$$

$$C_8 \rightarrow -\frac{9}{250}C_2^4 + \frac{4}{25}C_2C_3^2 + \frac{9}{25}C_2^2C_4 + \frac{1}{10}C_4^2 + \frac{4}{15}C_3C_5,$$

$$C_9 \rightarrow -\frac{13}{250}C_2^3C_3 + \frac{16}{225}C_3^3 + \frac{61}{150}C_2C_3C_4 + \frac{3}{10}C_2^2C_5 + \frac{1}{10}C_4C_5.$$

However, the non-primitive charges are not commuting. They are not Casimirs and we can not consider them as Hamiltonians. The similar method of construction chiral currents for $SO(2l+1) = B_l$, $SP(2l) = C_l$ groups was used by Evans et.al. on the base of symmetric invariant tensors of de Azcarraga et.al.. In the defining representation these groups generators corresponding algebras t_μ satisfy rules:

$$[t_\mu, t_\nu] = 2if_{\mu\nu}^\lambda t_\lambda, \text{Tr}(t_\mu t_\nu) = 2\delta_{\mu\nu}, t_\mu \eta = -\eta t_\mu^t,$$

where η is euclidean or symplectic structure.

The symmetric tensor structure constants for these groups they introduced through completely symmetrized product of three generators of corresponding algebras:

$$t_{(\mu} t_\nu t_{\lambda)} = v_{\mu\nu\lambda}^\rho t_\rho, \quad (8)$$

where $v_{\mu\nu\lambda\rho}$ is totally symmetric tensor. The basic invariant symmetric tensors have the form :

$$V_{\mu\nu}^{(2)} = \delta_{\mu\nu}, \quad V_{(\mu_1\mu_2\dots\mu_{2N-1}\mu_{2N})}^{(2N)} = v_{(\mu_1\mu_2\mu_3}^{\nu_1} v_{\mu_4\mu_5}^{\nu_2} \dots v_{\mu_{2N-2}\mu_{2N-1}\mu_{2N})}^{\nu_{2N-3}}, \quad N = 2, \dots, \infty.$$

The invariant chiral currents $J^{(2N)}$ coincide to the basis invariant chiral currents $V^{(2N)}$:

$$J^{(2N)} = 2V_{\mu_1\dots\mu_{2N}}^{(2N)} U^{\mu_1} \dots U^{\mu_{2N}}.$$

The commutation relations of invariant chiral currents are PBs of hydrodynamic type:

$$\{V^{(M)}(x), V^{(N)}(y)\} = -MN V^{(M+N-2)}(x) \frac{\partial}{\partial x} \delta(x-y) - \frac{MN(N-1)}{M+N-2} \frac{\partial V^{(M+N-2)}(x)}{\partial x} \delta(x-y).$$

The commuting charges of these invariant chiral currents are dynamical Casimir operators on $SO(2l+1)$, $SP(2l)$. The metric tensor of Riemann space of invariant chiral currents is the following:

$$g_{MN}(V(x)) = -MN(M+N-2)V^{(M+N-2)}(x).$$

The commutation relations coincide to commutation relations, which was obtained by Evans et.al. .

We used relations for new symmetric invariant tensors $V_{(\mu_1\dots\mu_{2N})}^{(2N,1)}$, which we obtained during calculation PB :

$$V_{(\mu_1\dots\mu_{10})}^{(10,1)} = v_{(\mu_1\mu_2\mu_3}^k v_{\mu_4\mu_5\mu_6}^l v_{\mu_7\mu_8\mu_9}^n v_{\mu_{10})}^{kl n} = V_{(\mu_1\dots\mu_{10})}^{(10)},$$

$$V_{(\mu_1 \dots \mu_{12})}^{(12,1)} = v_{(\mu_1 \mu_2 \mu_3)}^k v_{\mu_4 \mu_5 \mu_6}^l v_{\mu_7 \mu_8 \mu_9}^n v_{\mu_{10} \mu_{11} \mu_{12}}^m v^{klmn} = V_{(\mu_1 \dots \mu_{12})}^{(12)},$$

$$V_{(\mu_1 \dots \mu_{14})}^{(14,1)} = v_{(\mu_1 \mu_2 \mu_3)}^k v_{\mu_4 \mu_5 \mu_6}^l v_{\mu_7 \mu_8 \mu_9}^n v_{\mu_{10} \mu_{11} \mu_{12}}^m v_{\mu_{13} \mu_{14}}^{klp} v^{nmp} = V_{(\mu_1 \dots \mu_{14})}^{(14)}.$$

The invariant chiral currents $J^{(2N)}$ and $V^{(2N)}$ and the new dependent totally symmetric invariant tensors for $SO(2l+1)$, $SP(2l)$ groups can be obtained under different order of calculation of trace of the product of the generators of corresponding algebras. Let us mark the matrix product of three generators t_μ in round brackets:

$$(t_{(\mu} t_\nu t_{\lambda)}) = v_{\mu\nu\lambda\rho} t_\rho.$$

The different disposition of this triplet inside of J^{2N} produce different expressions for V^{2N} :

$$J^{(10)} = Tr[(\underline{ttt})t(\underline{ttt})(\underline{ttt})] = 2V^{(10)}, \quad J^{(10)} = Tr[(\underline{ttt})(\underline{ttt})(\underline{ttt})t] = 2V^{(10,1)},$$

$$J^{(12)} = Tr[t(\underline{ttt})t(\underline{ttt})t(\underline{ttt})] = 2V^{(12)}, \quad J^{(12)} = Tr[(\underline{ttt})(\underline{ttt})(\underline{ttt})(\underline{ttt})] = 2V^{(12,1)},$$

$$J^{(14)} = Tr[(\underline{ttt})t(\underline{ttt})t(\underline{ttt})(\underline{ttt})] = 2V^{(14)}, \quad J^{(14)} = Tr[(\underline{ttt})(\underline{ttt})(\underline{ttt})(\underline{ttt})t] = 2V^{(14,1)}.$$

New invariant chiral tensors do not led to new invariant chiral currents. We obtained following expressions for non-primitive invariant currents V_N for $SO(2l+1)$, $SO(2l)$, $SP(2l)$ groups:

$$l = 1 : V_{2N} \rightarrow V_2^N,$$

$$l = 2 : V_6 \rightarrow -\frac{1}{2}V_2^3 + \frac{3}{2}V_2V_4, \quad V_8 \rightarrow -\frac{1}{2}V_2^4 + V_2^2V_4 + \frac{1}{2}V_4^2, \quad V_{10} \rightarrow -\frac{1}{4}V_2^5 + \frac{5}{4}V_2V_4^2,$$

$$l = 3 : V_8 \rightarrow \frac{1}{6}V_2^4 - V_2^2V_4 + \frac{1}{2}V_4^2 + \frac{4}{3}V_2V_6, \quad V_{10} \rightarrow \frac{1}{6}V_2^5 - \frac{5}{2}(V_2^3V_4 - V_2^2V_6 - V_4V_6).$$

The non-primitive charges for $l \geq 2$ are not commuting also.

4. CONCLUSIONS

In present paper author used dynamical polynomial chiral invariant charges to construct new integrable equations. It is possible for $SU(3)$, $SO(3)$, $SP(2)$ groups only. The non-primitive charges for groups of bigger rank are not Casimirs.

This work was supported by Grant of RFFI-NASU 38/50-2008.

2. I. Ya. Aref'eva, P.P. Kulish, E.R. Nissimov and S.J. Pacheva, preprint LOMI E-I-1978, unpublished.
3. V.D. Gershun, Problems of atomic science and technology **3(1)** part 1 16 (2007).
4. V.D. Gershun, SIGMA **4** 041 (2008); nlin.SI/0805.0656.
5. J.M. Evans, M. Hassan, N.J. MacKay, A.J. Mountain, Nucl. Phys. **B561**, 385 (1999); hep-th/9902008.
6. J.A. de Azcarraga, A.J. Macfarlane, A.J. Mountain and J.C. Perez Bueno, Nucl. Phys. **B510** 657 (1998); physics/9706006.
7. A. Klein, J. Math. Phys. **4** 1283 (1963).
8. Sudbery A., Ph.D. Thesis, Cambridge Univ. 1970; J. Phys. **A23(15)** L705 (1990).